Author Index

AALBERSE R.C. *see* KONIECZNY A.
ABO T. *see* TOYABE S.
ACEITUNO E., CASTAÑÓN S., JIMÉNEZ C., SUBIRÁ D., DE GÓRGOLAS M., FERNÁNDEZ-GUERRERO M., ORTÍZ F. & GARCÍA R. Circulating immune complexes from HIV-1$^+$ patients induces apoptosis on normal lymphocytes, 317
ACKERMANN U. *see* KREUTZ M.
AGEMATSU K. *see* NAKAZAWA T.
AKBAR A.N. *see* CRAIGEN J.L.
AKBAR SK.MD.F. *see* KUROSE K.
ALI S. *see* DOUGLAS M.S.
ALKHATIB G., MURATA K. & RODER J. C. Cellular distribution of a natural killer cell tumour recognition-related surface antigen in purified human lymphocytes, 173
ALONSO G.F. *see* PALERMO M.S.
ALONSO L. *see* VARAS A.
ANCUTA P., FAHMI H., PONS J.-F., LE BLAY K. & CHABY R. Involvement of the membrane form of tumour necrosis factor-α in lipopolysaccharide-induced priming of mouse peritoneal macrophages for enhanced nitric oxide response to lipopolysaccharide, 259
ANDREASSI L. *see* DI RENZO M.
ANDREESEN R. *see* KREUTZ M.
ARAI H. *see* TODA S.
ASAHARA H. *see* KOBATA T.
ASANO M. *see* HAYASHI M.
AUTERI A. *see* DI RENZO M.
AYALA A. *see* MELDRUM D.R.

BADOVINAC V. *see* TRAJKOVIC V.
BANCROFT G. *see* SMITH D.
BANERJEE A. *see* MELDRUM D.R.
BARBACANE R.C. *see* CONTI P.
BARTH P. *see* STEINIGER B.
BEDFORD P.A. *see* MCCARTHY D.A.
BEHR C. *see* MONSO-HINARD C.
BESSLER W. *see* KREUTZ M.
BHAN A. *see* ISRAEL E.J.
BISSON S. *see* SCHEBESCH C.
BIZINKAUSKAS C.B. *see* KONIECZNY A.
BLACKSTOCK R. & CASADEVALL A. Presentation of cryptococcal capsular polysaccharide (GXM) on activated antigen-presenting cells inhibits the T-suppressor response and enhances delayed-type hypersensitivity and survival, 334
BLUMBERG R.S. *see* ISRAEL E.J.
BÖHMIG G.A., KRIEGER P.-M., SÄEMANN M.D., WENHARDT C., POHANKA E. & ZLABINGER G.J. n-Butyrate downregulates the stimulatory function of peripheral blood-derived antigen-presenting cells: a potential mechanism for modulating T-cell responses by short-chain fatty acids, 234
BOND J.F. *see* KONIECZNY A.
BOOG C.J.P. *see* WIJBERG O.L.C.
BOONSTRA R. *see* SCHILIZZI B.M.
BRANDTZAEG P. *see* HELGELAND L.
BRAUER A.W. *see* KONIECZNY A.
BRENNER B., WEINMANN S., GRASSMÉ H., LANG F., LINDERKAMP O. & GULBINS E. L-selectin activates JNK via src-like tyrosine kinases and the small G-protein Rac, 214
BRICKELL P.M. *see* FAULKNER L.
BROWN K.A. *see* MCCARTHY D.A.
BROWN K.L. *see* MABBOTT N.A.
BRUCE M.E. *see* MABBOTT N.A.
BUZA J., SILEGHEM M., GWAKISA P. & NAESSENS J. CD5$^+$B lymphocytes are the main source of antibodies reactive with non-parasite antigens in *Trypanosoma congolense*-infected cattle, 226

CAIN B.S. *see* MELDRUM D.R.
CAROL M.J. *see* MACHUGH N.D.
CASADEVALL A. *see* BLACKSTOCK R.
CASTAÑÓN S. *see* ACEITUNO E.
CATALDO I. *see* CONTI P.
CAULFIELD J.J, HAWRYLOWICZ C.M.,[1]KEMENY D.M. & LEE T.H. GM-CSF increases the ability of cultured macrophages to support autologous CD4$^+$T-cell proliferation in response to *Dermatophagoides pteronyssinus* and PPD antigen, 123
CHABY R. *see* ANCUTA P.
CHANG M.-d.Y. *see* KHALILI H.
CHUN W.H. *see* LEE K.H.
CINADER B. *see* GORCZYNSKI R.M.
CLEVELAND J.C. JR *see* MELDRUM D.R.
COHEN S.B.A., CRAWLEY J.B., KAHAN M.C., FELDMANN M. & FOXWELL B. M. J. Interleukin-10 rescues T cells from apoptotic cell death: association with an upregulation of Bcl-2, 1
CONTI P., REALE M., FELICIANI C., FRYDAS S., TRAKATELLIS M., PLACIDO F.C., CATALDO I., DI GIOACCHINO M. & BARBACANE R. C. Augmentation of monocyte chemotactic protein-1 (MCP-1) and mRNA transcript in chronic inflammatory states induced by potassium permanganate ($KMnO_4$) *in vivo*, 300
CRAIGEN J.L., YONG K.L., JORDAN N.J., MACCORMAC P., WESTWICK J., AKBAR A.N. & GRUNDY J. E. Human cytomegalovirus infection up-regulates interleukin-8 gene expression and stimulates neutrophil transendothelial migration, 138
CRAWLEY J.B. *see* COHEN S.B.A.
CROCKER P.R. *see* STEINIGER B.
CUMBERBATCH M., DEARMAN R.J. & KIMBER I. Langerhans cells require signals from both tumour necrosis factor-α and interleukin-1β for migration, 388
CUNNINGHAM C. *see* WARD F.J.

DAVIS W.C. *see* MACHUGH N.D.
DE ALOE G. *see* DI RENZO M.
DE GÓRGOLAS M. *see* ACEITUNO E.
DE LEIJ L.F.M.H. *see* SCHILIZZI B.M.
DEARMAN R.J. *see* CUMBERBATCH M.
DESHPANDE R. *see* KHALILI H.
DI GIOACCHINO M. *see* CONTI P.
DI RENZO M., RUBEGNI P., DE ALOE G., PAULESU L., PASQUI A.L., ANDREASSI L., AUTERI A. & FIMIANI M. Extracorporeal photochemotherapy restores Th1/Th2 imbalance in patients with early stage cutaneous T-cell lymphoma, 99
DLUGONSKA H. *see* REICHMANN G.
DOUGLAS M.S., ALI S., RIX D.A., ZHANG J.-G. & KIRBY J. A. Endothelial production of MCP-1: modulation by heparin and consequences for mononuclear cell activation, 512
DUBREMETZ J.F. *see* REICHMANN G.
DUMONDE D.C. *see* MCCARTHY D.A.

EHLERS S. *see* SMITH D.
EISHI Y. *see* WAKABAYASHI A.
EKMAN L. *see* KJERRULF M.
EL-GHORR A.A. & NORVAL M. The role of interleukin-4 in ultraviolet B light-induced immunosuppression, 26
EREMIN O. *see* PURASIRI P.
ERNST J.D. *see* LARSSON M.

FAHMI H. *see* ANCUTA P.
FAULKNER L., PATEL M., BRICKELL P.M. & KATZ D. R. The role of the Fgr tyrosine kinase in the control of the adhesive properties of U937 monoblastoid cells and their derivatives, 519

FELDMANN M. *see* COHEN S.B.A.
FELICIANI C. *see* CONTI P.
FERNÁNDEZ C. *see* SVERREMARK E.
FERNÁNDEZ-GUERRERO M. *see* ACEITUNO E.
FIMIANI M. *see* DI RENZO M.
FISCHER H.-G. *see* REICHMANN G.
FORSUM U. *see* LARSSON M.
FOXWELL B.M.J. *see* COHEN S.B.A.
FROHN C., SCHLENKE P. & KIRCHNER H. The repertoire of HLA–Cw-specific NK cell receptors CD158 a/b (EB6 and GL183) in individuals with different HLA phenotypes, 567
FRYDAS S. *see* CONTI P.
FUKUDA M. *see* TOYABE S.

GARCÍA R. *see* ACEITUNO E.
GELDER C.M. *see* TSITOURA D.C.
GITLITS V.M., SENTRY J.W., MATTHEW M.L.S.M., SMITH A.I. & TOH B.-H. Autoantibodies to evolutionarily conserved epitopes of enolase in a patient with discoid lupus erythematosus, 354
GLEESON P.A. *see* SCARFF K.J.
GOERDT S. *see* SCHEBESCH C.
GORAI I. *see* TODA S.
GORCZYNSKI R.M., CINADER B., RAMAKRISHNA V., TERZIOGLU E., WAELLI Th. & WESTPHAL O. An antibody specific for interleukin-6 reverses age-associated changes in spontaneous and induced cytokine production in mice, 20
GRASSMÉ H. *see* BRENNER B.
GRAU G.E. *see* MONSO-HINARD C.
GRDIC D. *see* KJERRULF M.
GRUNDY J.E. *see* CRAIGEN J.L.
GULBINS E. *see* BRENNER B.
GWAKISA P. *see* BUZA J.

HABU Y. *see* SEKI S.
HADZIC O. *see* TRAJKOVIC V.
HAKIJ N. *see* SCHEBESCH C.
HAMAJIMA K. *see* TODA S.
HAMANO S. *see* SONODA K.-H.
HAMMARSTRÖM L. *see* MOUSAVI M.
HÄNSCH H. *see* SMITH D.
HARKEN A.H. *see* MELDRUM D.R.
HARRIS W.J. *see* WARD F.J.
HARRISON L.C. *see* YAMASHITA K.
HARTNELL A. *see* STEINIGER B.
HASEGAWA A. *see* KUBO S.
HASHIMOTO K. *see* KUBO S.
HASHIMOTO W. *see* SEKI S.
HASSAN J., RAINSFORD E. & REEN D. J. Linkage of protein kinase C-β activation and intracellular interleukin-2 accumulation in human naive CD4 T cells, 465
HAUSCHILDT S. *see* KREUTZ M.
HAWRYLOWICZ C.M. *see* CAULFIELD J.J
HAYASHI M., TAKENOUCHI N., ASANO M., KATO M., TSURUMACHI T., SAITO T. & MORO I. The polymeric immunoglobulin receptor (secretory component) in a human intestinal epithelial cell line is up-regulated by interleukin-1, 220
HEEMSKERK M.H.M. *see* WIJBERG O.L.C.
HELGELAND L., BRANDTZAEG P., ROLSTAD B. & VAAGE J. T. Sequential development of intraepithelial γδ and αβ T lymphocytes expressing CD8αβ in neonatal rat intestine: requirement for the thymus, 447
HERBST B. *see* STEINIGER B.
HEYS S.D. *see* PURASIRI P.
HIRAI M. *see* SANO H.
HIRAIDE H. *see* SEKI S.
HIROSE S. *see* KOBATA T.
HOFFMEYER F., WITTE K. & SCHMIDT R. E. The high-affinity FcγRI on PMN: regulation of expression and signal transduction, 544
HOLMDAHL R. *see* ZHAO Y.-X.
HOSOKAWA J.-I. *see* KAMIO E.

IIAI T. *see* TOYABE S.
ISHIGURO N. *see* YAMADA E.
ISHII N. *see* TODA S.
ISOBE K. *see* YAMADA E.
ISOBE K.-I. *see* SANO H.
ISRAEL E.J., TAYLOR S., WU Z., MIZOGUCHI E., BLUMBERG R.S., BHAN A. & SIMISTER N. E. Expression of the neonatal Fc receptor, FcRn, on human intestinal epithelial cells, 69
ISSEKUTZ A.C. *see* SHANG X.-Z.
see also WALTER U.M.
ISTURIZ M.A. *see* PALERMO M.S.
IVANYI J. *see* ZHU X.
IWATA H. *see* YAMADA E.

JENKINSON E.J. *see* TCHILIAN E.Z.
JEONG J.-Y., KIM K.-U. & JUE D.-M. Tosylphenylalanine chloromethyl ketone inhibits TNF-α mRNA synthesis in the presence of activated NF-κB in raw 264.7 macrophages, 267
JIMÉNEZ C. *see* ACEITUNO E.
JIMÉNEZ E. *see* VARAS A.
JOHANSSON M., WARD M. & LYCKE N. B-cell-deficient mice develop complete immune protection against genital tract infection with *Chlamydia trachomatis*, 422
JORDAN N.J. *see* CRAIGEN J.L.
JUE D.-M. *see* JEONG J.-Y.
JUILLARD P. *see* MONSO-HINARD C.

KAHAN M.C. *see* COHEN S.B.A.
KAMIO E., SAKAMOTO Y., KIMURA T., TAKAKUWA H., SAKURAI M., HOSOKAWA J.-I., NAKAGAWA K., YOSHIDA F. & TAGAMI K. Increased natural killer resistance to cyclosporine A by continuous doses of dexamethasone in rats, 407
KANEDA K. *see* ONODERA S.
KASAIAN M.T. *see* KONIECZNY A.
KATO M. *see* HAYASHI M.
KATO T. *see* KOBATA T.
KATZ D.R. *see* FAULKNER L.
KAWAMURA T. *see* TOYABE S.
KAYAGAKI N. *see* KOBATA T.
KEMENY D.M. *see* CAULFIELD J.J
see also TSITOURA D.C.
KHALILI H., DESHPANDE R. & CHANG M.-d.Y. The defective antigen-presenting activity of murine fetal macrophage cell lines, 487
KHAR A. *see* LAKSHMI RAO K.
KIM K.-U. *see* JEONG J.-Y.
KIMBER I. *see* CUMBERBATCH M.
KIMURA T. *see* KAMIO E.
KIRBY J.A. *see* DOUGLAS M.S.
KIRCHNER H. *see* FROHN C.
KISHIHARA K. *see* YAMADA H.
KJERRULF M., GRDIC D., EKMAN L., SCHÖN K., VAJDY M. & LYCKE N. Y. Interferon-γ receptor-deficient mice exhibit impaired gut mucosal immune responses but intact oral tolerance, 60
KNIES J.E.G. *see* WARD F.J.
KNIGHT S.C. *see* MCCARTHY D.A.
KOBATA T., TAKASAKI K., ASAHARA H., MAI HONG N., MASUKO-HONGO K., KATO T., HIROSE S., SHIRAI T., KAYAGAKI N., YAGITA H., OKUMURA K. & NISHIOKA K. Apoptosis with $FasL^+$ cell infiltration in the periphery and thymus of corrected autoimmune mice, 206
KODELJA V. *see* SCHEBESCH C.
KONG Y.-Y. *see* YAMADA H.
KONIECZNY A., MORGENSTERN J.P., BIZINKAUSKAS C.B., LILLEY C.H., BRAUER A.W., BOND J.F., AALBERSE R.C., WALLNER B.P. & KASAIAN M. T. The major dog allergens, *Can f* 1 and *Can f* 2, are salivary lipocalin proteins: cloning and immunological characterization of the recombinant forms, 577
KRAAL G. *see* WOLVERS D.A.W.
KRAUSE S.W. *see* KREUTZ M.
KREUTZ M., ACKERMANN U., HAUSCHILDT S., KRAUSE S.W., RIEDEL D., BESSLER W. & ANDREESEN R. A comparative analysis of cytokine production and tolerance induction by bacterial lipopeptides, lipopolysaccharides and *Staphyloccocus aureus* in human monocytes, 396

KRIEGER P.-M. *see* BÖHMIG G.A.
KUBO M. *see* KUBO S.
KUBO S., HASEGAWA A., HASHIMOTO K., SHIMIZU C., KUBO M., TADA T. & NAKAYAMA T. CD4 regulates the efficiency of an endogenous superantigen-induced clonal detection of TCRVβ11$^+$ cells in the periphery, 437
KUROSE K., AKBAR SK.MD.F., YAMAMOTO K. & ONJI M. Production of antibody to hepatitis B surface antigen (anti-HBs) by murine hepatitis B virus carriers: neonatal tolerance versus antigen presentation by dendritic cells, 494
KUSAKABE K-I. *see* TODA S.

LAKSHMI RAO K., VARALAKSHMI CH., MUBARAK ALI A. & KHAR A. Administration of anti-IL-12 antibody *in vivo* inhibits rejection of a rat histiocytoma and suppresses cytokine response in a tumour-bearing host, 381
LAMB J.R. *see* TSITOURA D.C.
LANG F. *see* BRENNER B.
LARSSON M., MAJEED M., ERNST J.D., MAGNUSSON K.-E., STENDAHL O. & FORSUM U. Role of annexins in endocytosis of antigens in immature human dendritic cells, 501
LE BLAY K. *see* ANCUTA P.
LEE K.H., YOON M.S. & CHUN W. H. The effects of monoclonal antibodies against iC3b receptors in mice with experimentally induced disseminated candidiasis, 104
LEE T.H. *see* CAULFIELD J.J
LILLEY C.H. *see* KONIECZNY A.
LINDERKAMP O. *see* BRENNER B.
LOU J.N. *see* MONSO-HINARD C.
LYCKE N. *see* JOHANSSON M.
LYCKE N.Y. *see* KJERRULF M.

MABBOTT N.A., BROWN K.L., MANSON J. & BRUCE M.E. T-lymphocyte activation and the cellular form of the prion protein, 161
MACCORMAC P. *see* CRAIGEN J.L.
MACEY M.G. *see* MCCARTHY D.A.
MACHUGH N.D., MBURU J.K., CAROL M.J., WYATT C.R., ORDEN J.A. & DAVIS W. C. Identification of two distinct subsets of bovine γδ T cells with unique cell surface phenotype and tissue distribution, 340
MAGNUSSON K.-E. *see* LARSSON M.
MAI HONG N. *see* KOBATA T.
MAJEED M. *see* LARSSON M.
MAK T.W. *see* YAMADA H.
MAKINO H. *see* OHTA Y.
MANSON J. *see* MABBOTT N.A.
MARIANO M. *see* MOURA A.C.N.
MASUKO-HONGO K. *see* KOBATA T.
MATSUNO T. *see* ONODERA S.
MATSUZAKI G. *see* SONODA K.-H.
MATTHEW M.L.S.M. *see* GITLITS V.M.
MBURU J.K. *see* MACHUGH N.D.
MCCARTHY D.A., MACEY M.G., BEDFORD P.A., KNIGHT S.C., DUMONDE D.C. & BROWN K. A. Adhesion molecules are upregulated on dendritic cells isolated from human blood, 244
MCKECHNIE A. *see* PURASIRI P.
MEISEL R. *see* REICHMANN G.
MELDRUM D.R., CAIN B.S., CLEVELAND J.C. JR, MENG X., AYALA A., BANERJEE A. & HARKEN A. H. Adenosine decreases post-ischaemic cardiac TNF-α production: anti-inflammatory implications for preconditioning and transplantation, 472
MENG X. *see* MELDRUM D.R.
METCALFE D.D. *see* VLIAGOFTIS H.
MÉVÉLEC M.-N. *see* REICHMANN G.
MIYAISHI O. *see* YAMADA E.
MIZOGUCHI E. *see* ISRAEL E.J.
MODY C.H. *see* SYME R.M.
MONSO-HINARD C., LOU J.N., BEHR C., JUILLARD P. & GRAU G. E. Expression of major histocompatibility complex antigens on mouse brain microvascular endothelial cells in relation to susceptibility to cerebral malaria, 53
MORENO J. *see* VARAS A.
MORGENSTERN J.P. *see* KONIECZNY A.
MORO I. *see* HAYASHI M.
MOURA A.C.N. & MARIANO M. Lipids from *Mycobacterium leprae* cell wall suppress T-cell activation *in vivo* and *in vitro*, 429
MOUSAVI M., RABBANI H. & HAMMARSTRÖM L. Characterization of the bovine ε gene, 369
MUBARAK ALI A. *see* LAKSHMI RAO K.
MUKASA A. *see* SONODA K.-H.
MULLER C. *see* SCHEBESCH C.
MUÑOZ J.J. *see* VARAS A.
MURATA K. *see* ALKHATIB K.

NAESSENS J. *see* BUZA J.
NAKAGAWA K. *see* KAMIO E.
NAKAMURA K. *see* WAKABAYASHI A.
NAKAMURA T. *see* SONODA K.-H.
NAKASHIMA I. *see* SANO H.
see also YAMADA E.
NAKAYAMA T. *see* KUBO S.
NAKAZAWA T., AGEMATSU K. & YABUHARA A. Later development of Fas ligand-mediated cytotoxicity as compared with granule-mediated cytotoxicity during the maturation of natural killer cells, 180
NISHIHIRA J. *see* ONODERA S.
NISHIOKA K. *see* KOBATA T.
see also TODA S.
NOMOTO K. *see* SONODA K.-H.
see also YAMADA H.
NOMURA A. *see* SONODA K.-H.
NORVAL M. *see* EL-GHORR A.A.

OGASAWARA K. *see* SEKI S.
OHTA Y., YAMANE M., SOHDA T. & MAKINO H. TAK-603 selectively suppresses Th1-type cytokine production and inhibits the progression of adjuvant arthritis, 75
OKADA E. *see* TODA S.
OKUDA K. *see* TODA S.
OKUMURA K. *see* KOBATA T.
ONJI M. *see* KUROSE K.
ONODERA S., SUZUKI K., MATSUNO T., KANEDA K., TAKAGI M. & NISHIHIRA J. Macrophage migration inhibitory factor induces phagocytosis of foreign particles by macrophages in autocrine and paracrine fashion, 131
ORDEN J.A. *see* MACHUGH N.D.
ORFANOS C.E. *see* SCHEBESCH C.
ORTÍZ F. *see* ACEITUNO E.
OWEN J.J.T. *see* TCHILIAN E.Z.

PALERMO M.S., ROSA F.A., ALONSO G.F. & ISTURIZ M.A. Fcγ receptor-dependent clearance is enhanced following lipopolysaccharide *in vivo* treatment, 536
PARISH C.R. *see* YAMASHITA K.
PARKHOUSE R.M.E. *see* YANG H.
PASQUI A.L. *see* DI RENZO M.
PATEL M. *see* FAULKNER L.
PAULESU L. *see* DI RENZO M.
PETTITT J.M. *see* SCARFF K.J.
PLACIDO F.C. *see* CONTI P.
POHANKA E. *see* BÖHMIG G.A.
PONS J.-F. *see* ANCUTA P.
POPADIC D. *see* TRAJKOVIC V.
PURASIRI P., MCKECHNIE A., HEYS S.D. & EREMIN O. Modulation *in vitro* of human natural cytotoxicity, lymphocyte proliferative response to mitogens and cytokine production by essential fatty acids, 166

QADRI A. Identification of specific recognition molecules on murine mononuclear phagocytes and B lymphocytes for Vi capsular polysaccharide: modulation of MHC class II expression on stimulation with the polysaccharide, 146

RABBANI H. *see* MOUSAVI M.
RAINSFORD E. *see* HASSAN J.
RAMAKRISHNA V. *see* GORCZYNSKI R.M.
REALE M. *see* CONTI P.

REEN D.J. *see* HASSAN J.
REICHMANN G., STACHELHAUS S., MEISEL R., MÉVÉLEC M.-N., DUBREMETZ J.F., DLUGONSKA H. & FISCHER H.-G. Detection of a novel 40 000 MW excretory *Toxoplasma gondii* antigen by murine Th1 clone which induces toxoplasmacidal activity when exposed to infected macrophages, 284
RICH B. E. Autocrine expression of interleukin-7 rescues lymphoid expansion in interleukin-7-deficient mice, 374
RIEDEL D. *see* KREUTZ M.
RIX D.A. *see* DOUGLAS M.S.
RODER J.C. *see* ALKHATIB G.
ROLSTAD B. *see* HELGELAND L.
ROSA F.A. *see* PALERMO M.S.
RUBEGNI P. *see* DI RENZO M.

SÄEMANN M.D. *see* BÖHMIG G.A.
SAITO H. *see* SANO H.
SAITO T. *see* HAYASHI M.
SAKAMOTO Y. *see* KAMIO E.
SAKURAI M. *see* KAMIO E.
SANO H., HIRAI M., SAITO H., NAKASHIMA I. & ISOBE K.-I. A nitric oxide-releasing reagent, S-nitroso-N-acetylpenicillamine, enhances the expression of superoxide dismutases mRNA in murine macrophage cell line RAW264–7, 118
SATOH M. *see* SEKI S.
SCARFF K.J., PETTITT J.M., VAN DRIEL I.R., GLEESON P.A. & TOH B.-H. Immunization with gastric H^+/K^+-ATPase induces a reversible autoimmune gastritis, 91
SCHEBESCH C., KODELJA V., MULLER C., HAKIJ N., BISSON S., ORFANOS C.E. & GOERDT S. Alternatively activated macrophages actively inhibit proliferation of peripheral blood lymphocytes and $CD4^+$T cells *in vitro*, 478
SCHILIZZI B.M., BOONSTRA R., THE T.H. & DE LEIJ L.F.M.H. Effect of B-cell receptor engagement on CD40-stimulated B cells, 346
SCHLENKE P. *see* FROHN C.
SCHMIDT R.E. *see* HOFFMEYER F.
SCHÖN K. *see* KJERRULF M.
SEKI S., HASHIMOTO W., OGASAWARA K., SATOH M., WATANABE H., HABU Y., HIRAIDE H. & TAKEDA K. Antimetastatic effect of $NK1^+$T cells on experimental haematogenous tumour metastases in the liver and lungs of mice, 561
SENTRY J.W. *see* GITLITS V.M.
SHANG X.-Z. & ISSEKUTZ A.C. β_2 (CD18) and β_1 (CD29) integrin mechanisms in migration of human polymorphonuclear leucocytes and monocytes through lung fibroblast barriers: shared and distinct mechanisms, 527
SHIMIZU C. *see* KUBO S.
SHIRAI T. *see* KOBATA T.
SILEGHEM M. *see* BUZA J.
SIMISTER N.E. *see* ISRAEL E.J.
SMITH A.I. *see* GITLITS V.M.
SMITH D., HÄNSCH H., BANCROFT G. & EHLERS S. T-cell-independent granuloma formation in response to *Mycobacterium avium*: role of tumour necrosis factor-α and interferon-γ, 413
SOHDA T. *see* OHTA Y.
SONODA K.-H., MATSUZAKI G., NOMURA A., YAMADA H., HAMANO S., NAKAMURA T., MUKASA A. & NOMOTO K. Macrophages activated by *Listeria monocytogenes* induce organ-specific autoimmunity, 274
STACHELHAUS S. *see* REICHMANN G.
STAINES N.A. *see* WARD F.J.
STEINIGER B., BARTH P., HERBST B., HARTNELL A. & CROCKER P. R. The species-specific structure of microanatomical compartments in the human spleen: strongly sialoadhesin-positive macrophages occur in the perifollicular zone, but not in the marginal zone, 307
STENDAHL O. *see* LARSSON M.
STOJKOVIC M.M. *see* TRAJKOVIC V.
SUBIRÁ D. *see* ACEITUNO E.
SUZUKI K. *see* ONODERA S.
SUZUKI S. *see* TOYABE S.
SVERREMARK E. & FERNÁNDEZ C. Immunogenicity of bacterial carbohydrates: cholera toxin modulates the immune response against dextran B512, 153
SYME R.M., WOOD C.J., WONG H. & MODY C. H. Both $CD4^+$ and $CD8^+$ human lymphocytes are activated and proliferate in response to *Cryptococcus neoformans*, 194

TADA T. *see* KUBO S.
TAGAMI K. *see* KAMIO E.
TAKAGI M. *see* ONODERA S.
TAKAKUWA H. *see* KAMIO E.
TAKASAKI K. *see* KOBATA T.
TAKEDA K. *see* SEKI S.
TAKENOUCHI N. *see* HAYASHI M.
TAKEUCHI A. *see* YAMADA E.
TARKOWSKI A. *see* ZHAO Y.-X.
TAYLOR S. *see* ISRAEL E.J.
TCHILIAN E.Z., OWEN J.J.T. & JENKINSON E. J. Anti-α4 integrin antibody induces apoptosis in murine thymocytes and staphylococcal enterotoxin B-activated lymph node T cells, 321
TERZIOGLU E. *see* GORCZYNSKI R.M.
THE T.H. *see* SCHILIZZI B.M.
TODA S., ISHII N., OKADA E., KUSAKABE K-I., ARAI H., HAMAJIMA K., GORAI I., NISHIOKA K. & OKUDA K. HIV-1-specific cell-mediated immune responses induced by DNA vaccination were enhanced by mannan-coated liposomes and inhibited by anti-interferon-γ antibody, 111
TOH B.-H. *see* GITLITS V.M.
see also SCARFF K.J.
TOYABE S., IIAI T., FUKUDA M., KAWAMURA T., SUZUKI S., UCHIYAMA M. & ABO T. Identification of nicotinic acetylcholine receptors on lymphocytes in periphery as well as thymus in mice, 201
TRAJKOVIC V., BADOVINAC V., POPADIC D., HADZIC O. & STOJKOVIC M. M. Cell-specific effects of pentoxifylline on nitric oxide production and inducible nitric oxide synthase mRNA expression, 402
TRAKATELLIS M. *see* CONTI P.
TSITOURA D.C., GELDER C.M., KEMENY D.M. & LAMB J. R. Regulation of cytokine production by human Th0 cells following stimulation with peptide analogues: differential expression of TGF-β in activation and anergy, 10
TSURUMACHI T. *see* HAYASHI M.

UCHIYAMA M. *see* TOYABE S.

VAAGE J.T. *see* HELGELAND L.
VAJDY M. *see* KJERRULF M.
VAN DER CAMMEN M.J.F. *see* WOLVERS D.A.W.
VAN DRIEL I.R. *see* SCARFF K.J.
VAN ROOIJEN N. *see* WIJBERG O.L.C.
VARALAKSHMI CH. *see* LAKSHMI RAO K.
VARAS A., VICENTE A., JIMÉNEZ E., ALONSO L., MORENO J., MUÑOZ J.J. & ZAPATA A. G. Interleukin-7 treatment promotes the differentiation pathway of T-cell-receptor-αβ cells selectively to the $CD8^+$ cell lineage, 457
VENKATAPRASAD N. *see* ZHU X.
VICENTE A. *see* VARAS A.
VLIAGOFTIS H. & METCALFE D. D. Characterization of adhesive interactions between mast cells and laminin isoforms: evidence of a principal role for α6 integrin, 553
VORDERMEIER H.M. *see* ZHU X.

WAELLI Th. *see* GORCZYNSKI R.M.
WAKABAYASHI A., EISHI Y. & NAKAMURA K. Regulation of experimental autoimmune orchitis by the presence or absence of testicular antigens during immunological development in SCID mice reconstituted with fetal liver cells, 84
WALLNER B.P. *see* KONIECZNY A.
WALTER U.M. & ISSEKUTZ A. C. Role of E- and P-selectin in migration of monocytes and polymorphonuclear leucocyes to cytokine and chemoattractant-induced cutaneous inflammation in the rat, 290

WARD F.J., KNIES J.E.G., CUNNINGHAM C., HARRIS W.J. & STAINES N. A. Natural antibodies that react with V-region peptide epitopes of DNA-binding antibodies are made by mice with systemic lupus erythematosus as disease develops, 354
WARD M. *see* JOHANSSON M.
WARREN H.S. *see* YAMASHITA K.
WATANABE H. *see* SEKI S.
WEINMANN S. *see* BRENNER B.
WENHARDT C. *see* BÖHMIG G.A.
WESTPHAL O. *see* GORCZYNSKI R.M.
WESTWICK J. *see* CRAIGEN J.L.
WIJBERG O.L.C., HEEMSKERK M.H.M., BOOG C.J.P. & VAN ROOIJEN N. Role of spleen macrophages in innate and acquired immune responses against mouse hepatitis virus strain A59, 252
WITTE K. *see* HOFFMEYER F.
WOLVERS D.A.W., VAN DER CAMMEN M.J.F. & KRAAL G. Mucosal tolerance is associated with, but independent of, up-regulation of Th2 responses, 328
WONG H. *see* SYME R.M.
WOOD C.J. *see* SYME R.M.
WU Z. *see* ISRAEL E.J.
WYATT C.R. *see* MACHUGH N.D.

YABUHARA A. *see* NAKAZAWA T.
YAGITA H. *see* KOBATA T.
YAMADA E., ISHIGURO N., MIYAISHI O., TAKEUCHI A., NAKASHIMA I., IWATA H. & ISOBE K. Differential display analysis of murine collagen-induced arthritis: cloning of the cDNA-encoding murine ATPase inhibitor, 571
YAMADA H. *see* SONODA K.-H.
YAMADA H., KONG Y.-Y., KISHIHARA K., MAK T.W. & NOMOTO K. $p56^{lck}$ is not essential for the T-cell response to allo-MHC antigens, 33
YAMAMOTO K. *see* KUROSE K.
YAMANE M. *see* OHTA Y.
YAMASHITA K., PARISH C.R., WARREN H.S. & HARRISON L.C. A multimeric form of soluble recombinant sheep LFA-3(CD58) inhibits human T-cell proliferation, 39
YANG H. & PARKHOUSE R. M. E. Differential expression of CD8 epitopes amongst porcine CD8-positive functional lymphocyte subsets, 45
YONG K.L. *see* CRAIGEN J.L.
YOON M.S. *see* LEE K.H.
YOSHIDA F. *see* KAMIO E.

ZAPATA A.G. *see* VARAS A.
ZHANG J.-G. *see* DOUGLAS M.S.
ZHAO Y.-X., HOLMDAHL R. & TARKOWSKI A. Induction of selective anergy by toxic shock syndrome toxin-1 in $CD8^+$ T cells, 188
ZHU X., VENKATAPRASAD N., IVANYI J. & VORDERMEIER H. M. Vaccination with recombinant vaccinia viruses protects mice against *Mycobacterium tuberculosis* infection, 6
ZLABINGER G.J. *see* BÖHMIG G.A.

Subject Index

Adhesion molecules:
on dendritic cells from human blood, 244
Adjuvant arthritis, progression:
suppression by TAK-603, 75
Allergens, dog:
cloning and characterization, 577
Annexins:
role in endocytosis of antigens in immature human dendritic cells, 501
Anti-interleukin-6:
and age-related changes in cytokine production, 20
Anti-interleukin-12:
suppression of antitumour response, 381
Antigen-presenting activity, defective:
murine fetal macrophages, 487
Apoptosis:
in corrected autoimmune mice, 206
induced by circulating immune complexes from HIV-1-infected patients, 317
induction by anti-α4 integrin antibody, 321
Autoantibodies:
to epitopes of enolase, 362
Autoimmune gastritis:
induced by gastric ATPase, 91
Autoimmune orchitis:
in reconstituted SCID mice, 84

B-cell receptor engagement:
effect on CD40-stimulated B cells, 346
B-cell-deficient mice:
protection against *Chlamydia trachomatis* infection, 422

Candidiasis:
effects of monoclonal anti-iC3b receptors, 104
CD4:
regulation of superantigen-induced clonal deletion of TCRVβ11$^+$ cells, 437
CD5$^+$ B lymphocytes, bovine:
and non-specific antibodies, 226
CD8 epitopes:
porcine, 45
γδ cells, bovine:
identification of two subsets, 340
Cholera toxin:
modulation of immune response to dextran, 153
Collagen-induced arthritis:
differential display analysis, 571
Cryptococcis:
enhancement of delayed-type hypersensitivity and survival in, 334
Cytokine production:
by human Th0 cells, 10
in human monocytes, 396

Dexamethasone:
increase in natural killer resistance to cyclosporine A, 407
DNA vaccination, induction of HIV-1-specific responses:
enhancement by mannan-coated liposomes, 111

Essential fatty acids:
modulation of human cellular immune responses, 166
Excretory antigen, from *Toxoplasma gondii*:
stimulation of T-helper 1 clone, 284

Fcγ receptor-dependent clearance:
enhanced by lipopolysaccharide, 536
FcγRI, high affinity, on polymorphonuclear cells:
regulation of expression and signal transduction, 544
FcRn:
expression on human intestinal epithelial cells, 69

ε gene, bovine:
characterization, 369
Granulocyte macrophage–colony-stimulating factor:
effects on cultured macrophages, 123
Granuloma formation, T-cell independent:
in *Mycobacterium avium* infection, 413

Heparin:
modulation of cytokines, 512
Hepatitis B surface antigen, antibody to:
in murine hepatitis B virus carriers, 373

Immunosuppression:
by recombinant sheep LFA-3, 39
Interferon-γ receptor knockout mice:
mucosal immune response, 60
Interleukin-4:
role in ultraviolet B-induced immunosuppression, 26
Interleukin-7:
autocrine expression and lymphoid expansion, 374
promotion of differentiation of CD8$^+$ cells, 457
Interleukin-8:
up-regulation by human cytomegalovirus, 138
Interleukin-10:
rescue of T cells from apoptosis, 1

L-selectin:
stimulation of stress-activated protein kinases, 214
Langerhans cells, migration:
requirement for both tumour necrosis factor-α and interleukin-1β, 388
Lymphocytes, human, CD4$^+$ and CD8$^+$:
responses to *Cryptococcus neoformans*, 194

Macrophage migration inhibitory factor:
induction of phagocytosis by macrophages, 131
Macrophages, activated by *Listeria monocytogenes*:
induction of autoimmunity, 274
Macrophages, alternatively activated:
and lymphocyte proliferation, 478
Macrophages, splenic:
role in responses to mouse hepatitis virus, 252
Malaria, cerebral:
and expression of major histocompatibility complex antigen on brain endothelial cells, 53
Mast cell adhesion:
to laminin, 553
Migration across fibroblasts, polymorphonuclear leucocytes and monocytes:
integrin mechanisms, 527
Monocyte chemotactic protein:
augmentation in permanganate-induced inflammation, 300
Mucosal tolerance:
and T-helper 2 responses, 328

n-Butyrate:
modulation of T-cell responses, 234
Natural antibodies:
reacting with V-region epitopes, 354
Natural killer cell receptors, human:
human leucocyte antigen, 567

Natural killer cell tumour recognition factor:
in human lymphocytes, 173
Natural killer cells:
cytotoxicity during maturation, 180
Natural killer T cells:
anti-metastatic effect on tumour metastases, 561
Nicotinic acetylcholine receptors:
on lymphocytes, 201

Ontogeny of intraepithelial T cells in rat intestine, 447

$p56^{lck}$:
and T-cell response to allo–major histocompatibility complex antigens, 33
Pentoxifylline:
effects on nitric oxide production and nitric oxide synthase mRNA expression, 402
Prion protein, cellular form:
and T-cell activation, 161
Protein kinase C-β activation and intracellular interleukin-2 accumulation in naive $CD4^+$ T cells, 465

Recombinant vaccinia virus:
protection against *Mycobacterium tuberculosis*, 6
Secretory component:
up-regulation by interleukin-1, 220
Selectins:
role in leucocyte migration, 290
Sialoadhesin-positive macrophages:
in human spleen, 307
Superoxide dismutases, enhanced expression:
by S-nitroso-N-acetylpenicillamine, 118

T-cell activation:
suppression by *Mycobacterium leprae* lipids, 429
T-helper 1/T-helper 2 balance, in lymphoma:
restoration by photochemotherapy, 99
Toxic shock syndrome toxin-1:
induction of anergy in $CD8^+$ T cells, 188
Tumour necrosis factor-α, membrane form:
role in lipopolysaccharide priming of macrophages, 259
Tumour necrosis factor-α, RNA synthesis:
inhibition by chloromethyl ketones, 267
Tumour necrosis factor-α production, cardiac:
decreased by adenosine, 472
Tyrosine kinase:
role in adhesive properties of U937 cells, 519

Vi capsular polysaccharide:
recognition molecules on mononuclear phagocytes, 146